

TEMPORAL PROFILES AND SPECTRAL LAGS OF XRF 060218

EN-WEI LIANG^{1,2}, BIN-BIN ZHANG^{1,3,4}, MIKE STAMATIKOS⁵, BING ZHANG¹, JAY NORRIS⁵, NEIL GEHRELS⁵, JIN ZHANG^{3,4}, Z. G. DAI^{6,1}

¹Department of Physics, University of Nevada, Las Vegas, NV 89154, USA;
 lew@physics.unlv.edu; bzhang@physics.unlv.edu

²Department of Physics, Guangxi University, Nanning 530004, China

³National Astronomical Observatories/Yunnan Observatory, CAS, Kunming 650011, China

⁴The Graduate School of the Chinese Academy of Sciences, Beijing 100039, China

⁵NASA Goddard Space Flight Center, Greenbelt, MD 20771

⁶Department of Astronomy, Nanjing University, Nanjing 210093, China

Draft version February 5, 2008

ABSTRACT

The spectral and temporal properties of the non-thermal emission of the nearby XRF 060218 in 0.3-150 keV band are studied. We show that both the spectral energy distribution and the light curve properties suggest the same origin of the non-thermal emission detected by *Swift* BAT and XRT. This event has the longest pulse duration and spectral lag observed to date among the known GRBs. The pulse structure and its energy dependence are analogous to typical GRBs. By extrapolating the observed spectral lag to the *CGRO/BATSE* bands we find that the hypothesis that this event complies with the same luminosity-lag relation with bright GRBs cannot be ruled out at 2σ significance level. These intriguing facts, along with its compliance with the Amati-relation, indicate that XRF 060218 shares the similar radiation physics as typical GRBs.

Subject headings: gamma-rays: bursts—method: statistical

1. INTRODUCTION

X-ray flashes (XRFs), cosmic explosions with lower spectral peak energies (E_p) in νF_ν spectra than typical gamma-ray bursts (GRBs) (Heise et al. 2001; Kippen et al. 2003), are thought to be the low energy extension of typical GRBs (Lamb et al. 2005; Sakamoto et al. 2004, 2006; Cui et al. 2005). They may be GRB jets (uniform or structured) viewed at off-axis directions (e.g. Zhang et al. 2003; Yamazaki et al. 2004; Zhang et al. 2004) or intrinsically different events (Lamb et al. 2005; Soderberg et al. 2005). It is long speculated that long duration GRBs originate from a relativistic jet emerging from a collapsing massive star progenitor (Woosley et al. 1993; Paczynski 1998; MacFadyen et al. 1999; Zhang et al. 2003), and associations of core-collapsing supernovae (SNe) with long GRB afterglows have been spectroscopically identified in a number of systems, including GRB980425/SN1998bw (Galama et al. 1998), GRB030329/SN2003dh (Stanek et al. 2003; Hjorth et al. 2003), and GRB031203/SN 2003lw (Malesani et al. 2004). The firm spectroscopic association of the nearby XRF 060218 (at $z=0.0331$, Mirabal et al. 2006) detected by Swift (Campana et al. 2006) with the Type Ic SN 2006aj was established recently (Modjaz et al. 2006; Pian et al. 2006; Sollerman et al. 2006; Mirabal et al. 2006; Cobb et al. 2006; Soderberg et al. 2006). This suggests that the progenitors of both GRBs and XRFs are related to the death of massive stars and XRFs are the low energy extension of the more “standard” GRBs.

Some empirical relations have been discovered from typical GRBs. It is interesting to verify whether XRF 060218 satisfies these relations as an approach to access its “standardness”. It has been reported that XRF 060218 complies with the isotropic energy vs. spectral peak energy ($E_{iso} - E_p$) relation derived from long GRBs (Amati et al.

2002) and other XRFs (Amati et al. 2006). The pulses in GRB light curves usually display a fast-rise-exponential-decay (FRED) shape and it has been found that the pulse width is related to the energy of the observational band as $\omega \propto E^{-0.4}$ (Fenimore et al. 1995; Norris et al. 2005). A correlation between the isotropic luminosity and the spectral lag ($L_{iso} - \tau$ relation) of light curves was also discovered with typical GRBs as $L_{iso} \propto \tau^{-1.18}$ (Norris et al. 2000). Both $\omega - E$ and $L_{iso} - \tau$ relations are related to the structure of light curves. Since most photons of an XRF are in the X-ray band, previous GRB missions did not observe the real light curves of XRFs. This makes it difficult to identify the temporal structure and the spectral lag of XRFs. XRF 060218 is long and soft, and *Swift* BAT and XRT simultaneously collected the data in the gamma-ray to X-ray bands. This makes it possible to measure its temporal structure and to examine whether it complies with the same $\omega - E$ and $L_{iso} - \tau$ relations derived from GRBs. In this *Letter* we focus on this issue. Throughout the paper $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, and $\Omega_\Lambda = 0.7$ are adopted.

2. DATA

XRF 060218 was detected with the *Swift*/BAT on 2006 February 18.149 UT. It is a long burst, with a duration $T_{90} \sim 2000$ seconds in the 15-150 keV band. Swift slewed autonomously to the burst and the X-ray telescope (XRT) and UV/Optical Telescope (UVOT) began collecting data 159 s after the burst trigger.

XRF 060218 was an image trigger. The BAT event data lasted only until $t \sim 300$ seconds after the trigger. The survey data are used to derive the BAT light curve. XRF 060218 is very soft, with most of the emission in the BAT band being lower than 50 keV (Campana et al. 2006). In order to obtain a high level of signal-to-noise

ratio of the BAT light curve, we use the light curve in whole BAT band, i.e., 15-150 keV. The first orbit of the XRT data is fully in the Windowed Timing (WT) mode. We extract the light curves and spectrum of the XRT data with the Xselect package. The spectrum is grouped with the tool *grppha*, and the spectral fitting is carried out with the *Xspec* package. Following Campana et al. (2006) we fit the XRT spectrum in the first orbit with a model combining a black body component with temperature kT and a cut-off power law ($F \propto E^{-\Gamma} e^{-E/E_c}$) component. The absorption in both Milky Way¹ and the GRB host galaxy (N_H^{host}) is incorporated. We obtain $N_H^{host} = 0.63_{-0.03}^{+0.03} \times 10^{22} \text{ cm}^{-2}$, $kT = 0.122_{-0.004}^{+0.003} \text{ keV}$, $\Gamma = 1.78_{-0.08}^{+0.08}$, and $E_c = 21.8_{-6.7}^{+15.8} \text{ keV}$ (corresponding to $E_p \sim 5 \text{ keV}$), with reduced $\chi^2 = 1.55$ for 769 degrees of freedom. We derive the unabsorbed light curves in the 0.3-2 keV, 2-5 keV, and 5-10 keV bands from the XRT data. As reported by Campana et al. (2006), a thermal component exists, and is likely of different origin (e.g. the shock breakout emission associated with SN 2006aj; c.f. Li 2006; Ghisellini et al. 2006b) from the non-thermal component. We therefore subtract the contribution of this component from the observed light curves. The kT and the radiation radii (R_{BB}) evolve with time during the first orbit. We read the values of kT and R_{BB} from Campana et al. (2006) and calculate the light curves of this component in 0.3-2, 2-5, and 5-10 keV bands. Since the temperature of the black body component is below 0.2 keV, the light curves in the energy band higher than 2 keV is essentially not contaminated by the thermal component. In the 0.3-2 keV band, the derived light curve of the thermal component continuously increase with time, which could be well fitted by $\log F_{0.3-2} = (-10.69 \pm 0.09) + (0.66 \pm 0.03) \log t$ in cgs units. We thus subtract the contribution of this component from the 0.3-2 keV band light curve according to this fitting result. The derived non-thermal light curves in the three XRT bands as well as the one in the BAT band are shown in Fig.1(a). To clearly view the pulse width and spectral lag dependences with energy, we also plot in Fig.1(b) the normalized lightcurves. The characteristics of the light curves along with the average photon energy ($\bar{E}$) of each energy band are reported in Table 1.

3. RESULTS

3.1. Spectral energy distribution

One important question is whether the XRT non-thermal emission and the BAT emission are of the same origin. In order to clarify this we first study the spectral energy distribution (SED) of the burst using joint BAT-XRT data. Since this event is image trigger, the BAT event file contains only the data in the first 300 seconds since the BAT trigger. XRT began collecting data 159 seconds after the BAT trigger. We thus only obtain simultaneous observations of the two instruments from 160 to 300 seconds with the event files. The joint-fit SED in this period is shown in Fig.2, which is well fitted by the BB+CPL model (Campana et al. 2006) with the following parameters: $N_H^{host} = 0.65_{-0.03}^{+0.03} \times 10^{22} \text{ cm}^{-2}$, $kT = 0.13_{-0.01}^{+0.01} \text{ keV}$, $\Gamma = 1.56_{-0.07}^{+0.08}$ and $E_c = 122_{-46}^{+160} \text{ keV}$ (corresponding to $E_p \sim 54 \text{ keV}$), with reduced $\chi^2 = 0.97$ for 327 degrees of

freedom. The E_p strongly evolves with time, from 54 keV at the beginning down to $\sim 5 \text{ keV}$ at later times. This is consistent with that reported by Campana et al. (2006) and Ghisellini et al. (2006a). The model fitting results are shown in Fig.2. One can observe that the BAT-component is a good extrapolation of the XRT non-thermal component. This result implies that the non-thermal emissions detected by XRT and BAT are of the same origin.

3.2. Pulse width and Energy Dependence

As shown in Fig.1, the light curves in different energy bands can be all modeled by a single FRED-pulse. Koccevski et al. (2003) developed an empirical expression to fit a FRED-like pulse, which reads,

$$F(t) = F_m \left(\frac{t + t_0}{t_m + t_0} \right)^r \left[\frac{d}{d+r} + \frac{r}{d+r} \left(\frac{t + t_0}{t_m + t_0} \right)^{(r+1)} \right]^{-\frac{r+d}{r+1}}, \quad (1)$$

where t_m is the time of the maximum flux (F_m), t_0 is the offset time, and r and d are the rising and decaying power-law indices, respectively. We fit the light curves with Eq.(1) and then measure the pulse width, rising and decaying times at the full-width half-maximum (FWHM) of the fitting light curves, and the rising-to-decaying time ratio (φ). The errors of these quantities are derived from simulations by assuming a normal distribution of the errors of the fitting parameters. The reported errors are at 1σ confidence level. The results are tabulated in Table 1. We show ω as a function of $\bar{E}$ in Fig.3(a). Apparently the two quantities are correlated. A best fit yields $\omega \propto E^{-0.31 \pm 0.03}$. The φ parameter ranges from 0.43 to 0.59. It is found that XRF 060218 roughly satisfies the same $\omega - E$ relation (Fenimore et al. 1995; Norris et al. 2005) and its φ values are also well consistent with that observed in typical GRBs (e.g. Norris et al. 1996; Liang et al. 2002), although it has a much longer pulse width than other single-pulse GRBs. These results imply that XRF 060218 may be an extension of GRBs to the extremely long and soft regime.

3.3. Spectral Lag and Energy Dependence

The light curves shown in Fig.1(a) display a significant spectral lag (τ), with soft photons lagging behind the hard photons, as usually seen in long GRBs (Norris et al. 2000; Yi et al. 2006). We illustrate this lag behavior with the intensity-normalized light curves in Fig.1(b). The light curves peak at 405 ± 25 , 735 ± 9 , 919 ± 7 , and 1082 ± 13 seconds, respectively, in a sequence of high energy band to low energy band as shown in Fig. 3(b). The best fit to the correlation between the peak time (t_{peak}) and the average photon energy yields,

$$\log t_{peak} = (3.04 \pm 0.04) - (0.25 \pm 0.05) \log E. \quad (2)$$

A simple estimate of the lags between any pairs of the 4 light curves obtains $\tau_{peak} = 163 \sim 677$ seconds, being consistent with that shown in Gehrels et al. (2007). We note XRF 060218 becomes the new record-holder of the long-lag, wide-pulse GRBs. The previous record holder was GRB 971208, with $\tau \sim 58$ seconds and $\omega = 395$ seconds (Norris et al. 2005).

¹The N_H of our Galaxy (Dickey et al. 1990) for this burst is $\sim 1.1 \times 10^{21} \text{ cm}^{-2}$

We also calculate the lags with the cross correlation function (CCF) method. The errors of lags are evaluated by simulations. The results are also reported in Table 1. The lag derived by the CCF method (τ_{CCF}) is strongly correlated with τ_{peak} , but is systematically lower² than τ_{peak} [Fig. 4(a)]. A best fit gives $\tau_{\text{CCF}} = (-100 \pm 17) + (0.91 \pm 0.08)\tau_{\text{peak}}$.

The $L_{\text{iso}} - \tau$ relation was discovered with six bright BATSE GRBs (Norris et al. 2000) and the spectral lag was defined by the light curves in the 25-50 keV and 100-300 keV bands. We investigate whether the lag behavior of XRF 060218 is consistent with the $L_{\text{iso}} - \tau$ relation. Since XRF 060218 is a soft XRF and the emission in the 100-300 keV band is too weak to derive a light curve, we assume that t_{peak} of the light curve in the 100-300 keV band follows the $t_{\text{peak}} - E$ relation (Eq.2) and perform the extrapolation. With the extrapolated t_{peak} we then estimate τ_{peak} for the light curves in the 25-50 keV (average energy 30 keV) and 100-300 keV (average energy 200 keV) bands. We obtain $\tau_{\text{peak}} = 177 \pm 16$ seconds. Since τ_{CCF} is more reliable, we use the $\tau_{\text{peak}} - \tau_{\text{CCF}}$ relation [Fig. 4(a)] to derive $\tau_{\text{CCF}} = 61 \pm 26$ seconds. This lag is used in the $L - \tau$ relation analysis. Using the peak fluxes in the BAT and XRT band, we estimate $L_{\text{iso}} = 1.2 \times 10^{47}$ ergs s⁻¹.

Figure 4(b) shows the $L_{\text{iso}} - \tau$ relation derived by Norris et al. (2000) compared against XRF 060218 as well as two other nearby GRBs, 980425 and 031203. The data of the previous GRBs are taken from Norris et al. (2000) and Sazonov et al. (2004)³. The grey band and the two dashed lines mark the best fits at the 1σ and 2σ confidence level, respectively, and the solid line is the regression line for the 6 GRBs that were used to draw the $L - \tau$ correlation, i.e. $\log L_{\text{iso}} = (50.22 \pm 0.32) - (1.21 \pm 0.21) \times \log \tau$ (errors are at the 1σ level). We can see that XRF 060218 is definitely inside the 2σ region and is marginally at the 1σ region boundary. Therefore, the hypothesis that XRF 060218 follows the $L - \tau$ relation cannot be ruled out at the 2σ significance level. We caution that the τ is inferred from the extrapolation of the $t_{\text{peak}} - E$ relation. This introduces uncertainties in deriving the lag. The other two nearby GRBs, 980425 and 031203, are out of the 2σ region, which are identified as significant outliers of this relation (e.g. Sazonov et al. 2004).

4. CONCLUSIONS AND DISCUSSION

We have investigated the non-thermal emission of XRF 060218. The early SED of this event from 0.3-150 keV observed by BAT and XRT suggests that the non-thermal emission detected by the two instruments are the same component. By subtracting the contribution of the thermal emission we derive the light curves of the non-thermal emission. They are composed of a broad single pulse, and

the energy dependences of the widths and the rising-to-decaying-time ratio of the pulses are roughly consistent with those derived in typical GRBs. The light curves show significant spectral lags, with a well-defined peak time sequence from high energy band to low energy bands, i.e. $t_{\text{peak}} \propto E^{-0.25 \pm 0.05}$. We infer the spectral lag in the BATSE bands and find that the hypothesis that this event complies with the $L_{\text{iso}} - \tau$ relation with typical GRBs cannot be ruled out at the 2σ significance level.

These intriguing facts, along with its compliance with the Amati-relation, strongly suggest that GRB 060218 is a “standard” burst at the very faint, long, and soft end of the GRB distribution. Since all these relations concern the temporal and spectral properties of emission, they are likely related to the radiation mechanisms. The results therefore imply that XRF 060218 and other XRFs may share the similar radiation physics (e.g. synchrotron or inverse Compton scattering in internal shocks, Mészáros, 2002; Zhang & Mészáros 2004; Piran 2005; Mészáros 2006) with harder GRBs.

As discovered by Norris (2002), the proportion of long-lag bursts within long-duration bursts increases from negligible among bright BATSE bursts to $\sim 50\%$ at the trigger threshold, and their peak fluxes are ~ 2 orders of magnitude lower than those of the brightest bursts. This argues that they are intrinsically under-luminous. Taken together with the fact that three nearby GRBs, 980425, 031203, and 060218, are long-lagged and under-luminous, an intuitive speculation is that long-lag bursts are probably relatively nearby (e.g., Norris et al. 2005). The local GRB rate of these GRBs thus should be much higher than that expected from the high luminosity GRBs (Liang et al. 2006; see also Cobb et al. 2006; Pian et al. 2006; Soderberg et al. 2006). A possible scenario to explain their wide-pulse, long-lag, and under-luminous features is the off-axis viewing angle effect (e.g. Nakamura 1999; Salmonson 2000; Ioka & Nakamura 2001). Another scenario is that these features are intrinsic, being due to their lower Lorentz factors (Kulkarni et al. 1998; Woosley & MacFadyen 1999; Salmonson 2000; Dai et al. 2006; Wang et al. 2006). They might be from a unique GRB population (Liang et al. 2006) having a different type of central engine (e.g. neutron stars rather than black holes) from bright GRBs (e.g. Mazzali et al. 2006; Soderberg et al. 2006).

We thank the anonymous referee for helpful suggestions, and S. Campana, D. Burrows, J. Nousek, K. Page, T. Sakamoto, X.-Y. Wang, and Z. Li for discussion. This work was supported by NASA under grants NNG06GH62G and NNG05GB67G, and the National Natural Science Foundation of China under grants 10463001(EWL).

²We here derive τ_{CCF} from the peak of the CCF without considering the side lobe contribution of the CCF. A fit to the CCF with a cube or quartic function gives a larger lag by considering the side lobe contribution (Norris et al. 2000). However, this method strongly depends on the artificially-selected range of CCF for the fitting. Since the light curves are a smooth pulse and their lags are significantly larger than the time bin, the peaks of CCFs are robust to estimate the lags.

³The peak photon fluxed in the 50-300 keV bands of these GRBs are converted to energy fluxes with the spectral index presented in Friedman & Bloom (2005), and their L_{iso} 's are recalculated with the cosmological parameters used in this paper.

REFERENCES

- Amati, L., et al., 2002, *A&A*, 390, 81
 Amati, L., et al., 2006, *GCN* 4846.
 Campana, S., et al. 2006, *Nature*, 442, 1008
 Cobb, B. E., et al. 2006, *ApJ*, 645, L113
 Cui, X. H., Liang, E. W., & Lu R. J., 2005, *ChJAA*, 5, 151
 Dai, Z. G., Zhang, B., & Liang, E. W. 2006, *astro-ph/0604510*
 Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
 Fenimore, E. E., et al. 1995, *ApJ*, 448, L101
 Friedman, A. S., & Bloom, J. S. 2005, *ApJ*, 627, 1
 Galama, T. J. et al. 1998, *Nature*, 395, 670
 Gehrels, G., et al. 2006, *Nature*, in press (*astro-ph/0610635*).
 Ghisellini, G., et al. 2006a, *MNRAS*, in press (*astro-ph/0605431*)
 Ghisellini, G., Ghirlanda, G., Tavecchio, F. 2006b, *MNARS*, submitted (*astro-ph/0608555*)
 Heise, J., et al. 2001, in *Gamma-Ray Bursts in the Afterglow Era*, ed. E. Costa, F. Frontera, & J. Hjorth (Berlin: Springer), 16
 Hjorth, J., et al. 2003, *Nature*, 423, 847
 Ioka, K., Nakamura, T. 2001, *ApJ*, 554, L163
 Kippen, R. M., et al. 2003, in *AIP Conf. Proc.* 662, *Gamma-Ray Burst and Afterglow Astronomy 2001*, ed. G. R. Ricker & R. K. Vanderspek (Melville: AIP), 244
 Kocevski, D., Ryde, F., and Liang, E. 2003, *ApJ*, 596, 389
 Kulkarni, S. R., et al. 1998, *Nature*, 395, 663
 Li, L. X. 2006, *MNARS*, submitted (*astro-ph/0605387*)
 Liang, E. W., Xie, G. Z., Su, C. Y. 2002, *PASJ*, 54, 1
 Liang, E. W., Zhang, B., Virgili, F. & Dai, Z. G. 2006, *astro-ph/0605200*
 Lamb, D. Q., Donaghy, T. Q., Graziani, C. 2005, *ApJ*, 620, 355
 Mazzali, P. A. et al. 2006, *Nature*, 442, 1011
 Mészáros, P. 2002, *ARA&A*, 40, 137
 Mészáros, P. 2006, *Rep. Prog. Phys.*, 69, 2259
 MacFadyen, A. I. & Woosley, S. E. 1999, *ApJ*, 524, 262
 Malesani et al. 2004, *ApJ*, 609, L5
 Nakamura, T. 1999, *ApJ*, 522, L101
 Mirabal, N., et al. 2006, *ApJ*, 643, L99
 Modjaz, M., et al. 2006, *ApJ*, 645, L21
 Norris, J. P. et al. 1996, *ApJ*, 459, 393
 Norris, J. P., 2002, *ApJ*, 579, 386
 Norris, J. P. et al. 2005, *ApJ*, 627, 324
 Norris, J. P., Marani, G. F., Bonnell, J. T. 2000, *ApJ*, 534, 248
 Paczyński, B. 1998, *ApJ*, 494, L45
 Pian, E., et al. 2006, *Nature*, 442, 1011
 Piran, T. 2005, *Rev. Mod. Phys.*, 76, 1143
 Sakamoto, T., et al. 2004, *ApJ*, 602, 875
 Sakamoto, T., et al. 2006, *ApJ*, 636, L73
 Salmonson, J. D. 2000, *ApJ*, 544, L115
 Sazonov, S. Yu., Lutovinov, A. A., Sunyaev, R. A., 2004, *Nature*, 430, 646
 Soderberg, A. M., et al. 2005, *ApJ*, 627, 877
 Soderberg, A. M., et al. 2006, *Nature*, 442, 1014
 Sollerman, J., et al. 2006, *A&A*, 454, 503
 Stanek, K. Z., et al. 2003, *ApJ*, 591, L17
 Wang, X.-Y., Li, Z., Waxman, E. & Mészáros, P. 2006, *astro-ph/0608033*
 Woosley, S. E., et al. 1993, *ApJ*, 405, L273
 Woosley, S. E., & MacFadyen, A. I. 1999, *A&AS*, 138, 499
 Yamazaki, R., Ioka, K., Nakamura, T. 2004, *ApJ*, 607, L103
 Yi, T. F., Liang, E. W., Qin, Y. P., Lu, R. J. 2006, *MNRAS*, 367, 1751
 Zhang, B. & Mészáros, P., 2004, *Int. J. Mod. Phys. A*, 19, 2385
 Zhang, B. et al. 2004, *ApJ*, 601, L119
 Zhang, W., Woosley, S. E., & MacFadyen, A. I. 2003, *ApJ*, 586, 356

TABLE 1
TEMPORAL STRUCTURES AND SPECTRAL LAGS OF THE LIGHT CURVES

Temporal Structures

Band	Peak(s)	ω (s)	Rising time (s)	Decaying time (s)	φ	E (keV)
(1)15-150 keV	405(25)	889 (244)	311 (28)	578(185)	0.54(0.18)	36.9
(2)5-10 keV	735(9)	1278 (45)	475 (12)	803(35)	0.59(0.03)	6.9
(3)2-5 keV	919(7)	1707 (40)	624 (8)	1084(34)	0.58(0.02)	3.1
(4)0.3-2 keV	1082(13)	2625 (125)	794 (14)	1831 (112)	0.43(0.03)	0.7

Spectral Lags

Bands	ΔE (keV)	τ_{peak} (s)	τ_{CCF} (s)	Bands	ΔE (keV)	τ_{peak} (s)	τ_{CCF} (s)
(1)-(2)	30.0	330(26)	249(37)	(1)-(3)	33.8	514(26)	389(47)
(1)-(4)	36.2	677(28)	518(70)	(2)-(3)	3.8	184(11)	81(12)
(2)-(4)	6.2	347(16)	173(25)	(3)-(4)	2.4	163(15)	43(8)

Note: all the errors are derived by simulations and in 1σ significance level.




FIG. 1.— (a) Unabsorbed light curves of the non-thermal gamma-rays/X-rays in the energy bands of 15-150 keV, 5-10 keV, 2-5 keV, and 0.3-2 keV, respectively. The fitting curves with Eq.(1) are plotted. (b) Normalized light curves from the empirical model fitting.




FIG. 2.— The BAT-XRT joint spectral energy distribution from 160 to 300 seconds since the BAT trigger. The BB+CutoffPL fitting model is also shown.




FIG. 3.— The pulse duration (panel a) and the peak time (panel b) as a function of the average photon energy of the non-thermal emission. The solid lines in both panels are the best fits.




FIG. 4.— Panel (a): Comparison of the spectral lags derived from the peak times and from the CCF method. The solid line is the best fit. Panel (b): Isotropic gamma-ray luminosity as a function of spectral lag. The spectral lags of typical GRBs and GRB 980425 are calculated with the light curves in the 25-50 keV and 100-300 keV bands observed by CGRO/BATSE. The lag of GRB 031203 is calculated with the light curves in the 20-50 keV and 100-200 keV bands. The grey band and the two dashed lines mark the best fits at the 1 σ and 2 σ confidence level, respectively, and the solid line is the regression line for the six typical GRBs presented in Norris et al. (2000).